# © Copyright 2000

Gabriel Andrés Vecchi

## Sub-seasonal wind variability and El Niño

by

Gabriel Andrés Vecchi

A dissertation submitted in partial fulfillment of the requirements for the degree of

Doctor of Philosophy

University of Washington

2000

Program Authorized to Offer Degree: School of Oceanography

## University of Washington Graduate School

This is to certify that I have examined this copy of a doctoral dissertation by

### Gabriel Andrés Vecchi

and have found that it is complete and satisfactory in all respects, and that any and all revisions required by the final examining committee have been made.

Chair of Supervisory Committee:	
Don E. Harrison	
Reading Committee:	
Mitsuhiro Kawase	
Luanne Thompson	
Date:	

## University of Washington Graduate School

This is to certify that I have examined this copy of a doctoral dissertation by

### Gabriel Andrés Vecchi

and have found that it is complete and satisfactory in all respects, and that any and all revisions required by the final examining committee have been made.

Chair of Supervisory Committee:	
Don E. Harrison	-
Reading Committee:	
Mitsuhiro Kawase	
Luanne Thompson	
Date:	

#### **Doctoral Dissertation**

In presenting this dissertation in partial fulfillment of the requirements for the Doctoral degree at the University of Washington, I agree that the Library shall make its copies freely available for inspection. I further agree that extensive copying of the dissertation is allowable only for scholarly purposes, consistent with "fair use" as prescribed in the U.S. Copyright Law. Requests for copying and reproduction of this dissertation may be referred to Bell and Howell Information and Learning, 300 North Zeed Road, P.O. Box 1346, Ann Arbor, MI 48106-1346, or to the author.

Signature			
Č			
Date			

## University of Washington

#### Abstract

Sub-seasonal wind variability and El Niño

by Gabriel Andrés Vecchi

Chairperson of the Supervisory Committee

Affiliate Professor D.E. Harrison School of Oceanography

The role of sub-seasonal western and central Pacific surface zonal wind variability in the evolution of El Niño sea surface temperature anomaly (SSTA) is examined. The surface wind structure and atmospheric convection variability of tropical Pacific westerly wind events (WWEs) is described. The average tropical Pacific SSTA variability following and in the absence of equatorial WWEs is examined over the period 1986-1998. The surface wind structure of the Madden-Julian Oscillation is examined (MJO), along with relationships between the MJO and WWEs. The response of an ocean general circulation model (OGCM) to composite MJO and WWE surface wind stress forcing is explored.

WWEs can be classified into eight types based on the location of the maximum zonal wind anomalies. WWEs are found to be compact in space and time, exhibiting little translation during their lifetime. Typical scales for WWEs are derived over the period 1986-1995. It is found that certain WWE types exhibit significant seasonality, and some have a significant correlation with the Troup Southern Oscillation Index.

In the absence of equatorial WWEs, tropical Pacific SSTA tends to remain at or return towards climatology. Following equatorial WWEs, tropical Pacific SSTA warms towards or remains at El Niño type conditions.

WWEs are significantly associated with atmospheric convection. The MJO surface wind stress anomaly fields exhibit two regimes in the equatorial Pacific. In the western equatorial Pacific there are both easterly and westerly anomalies with the westerlies dominating, in the central and eastern equatorial Pacific the main anomalies are easterlies. The WWEs which occur during the MJO are modulated by the convective variability of the MJO. West-of-dateline WWEs exhibit a strong relationship to tropical cyclone activity.

Following a composite MJO, the tropical Pacific SSTA in the OGCM does not tend to warm; the composite MJO does not provide a simple mechanism for El Niño waveguide warming. Idealized WWEs drove tropical Pacific SSTA warming in the OGCM; enhanced WWE activity provides a mechanism for the onset of El Niño.